

## Contents

<b>1</b>	<b>Basic definitions</b>	<b>February 25, 2019</b>	<b>1</b>
1.1	Reidemeister moves . . . . .		3
1.2	Seifert surface . . . . .		4
1.3	Seifert matrix . . . . .		7
<b>2</b>		<b>March 4, 2019</b>	<b>8</b>
<b>3</b>			<b>11</b>
<b>4</b>	<b>Concordance group</b>	<b>March 18, 2019</b>	<b>12</b>
<b>5</b>		<b>April 8, 2019</b>	<b>14</b>
<b>6</b>		<b>March 11, 2019</b>	<b>15</b>
<b>7</b>		<b>April 15, 2019</b>	<b>15</b>
<b>8</b>		<b>May 20, 2019</b>	<b>16</b>
<b>9</b>		<b>May 27, 2019</b>	<b>22</b>
<b>10</b>		<b>June 3, 2019</b>	<b>23</b>
<b>11</b>	<b>balagan</b>		<b>24</b>

**Lecture 1 Basic definitions February 25, 2019**





### **Definition 1.1**

*A knot  $K$  in  $S^3$  is a smooth (PL - smooth) embedding of a circle  $S^1$  in  $S^3$ :*

$$\varphi : S^1 \hookrightarrow S^3$$

Usually we think about a knot as an image of an embedding:  $K = \varphi(S^1)$ .

**Example 1.1**

- *Knots:*  (*unknot*),  (*trefoil*).
- *Not knots:*  (*it is not an injection*),  (*it is not smooth*).

**Definition 1.2**

Two knots  $K_0 = \varphi_0(S^1)$ ,  $K_1 = \varphi_1(S^1)$  are equivalent if the embeddings  $\varphi_0$  and  $\varphi_1$  are isotopic, that is there exists a continuous function

$$\begin{aligned} \Phi : S^1 \times [0, 1] &\hookrightarrow S^3 \\ \Phi(x, t) &= \Phi_t(x) \end{aligned}$$

such that  $\Phi_t$  is an embedding for any  $t \in [0, 1]$ ,  $\Phi_0 = \varphi_0$  and  $\Phi_1 = \varphi_1$ .

**Theorem 1.1**

Two knots  $K_0$  and  $K_1$  are isotopic if and only if they are ambient isotopic, i.e. there exists a family of self-diffeomorphisms  $\Psi = \{\psi_t : t \in [0, 1]\}$  such that:

$$\begin{aligned} \psi(t) &= \psi_t \text{ is continuous on } t \in [0, 1] \\ \psi_t &: S^3 \hookrightarrow S^3, \\ \psi_0 &= id, \\ \psi_1(K_0) &= K_1. \end{aligned}$$

**Definition 1.3**





A knot is trivial (*unknot*) if it is equivalent to an embedding  $\varphi(t) = (\cos t, \sin t, 0)$ , where  $t \in [0, 2\pi]$  is a parametrisation of  $S^1$ .

**Definition 1.4**

A link with  $k$  - components is a (smooth) embedding of  $\overbrace{S^1 \sqcup \dots \sqcup S^1}^k$  in  $S^3$




**Example 1.2**

*Links:*

- a trivial link with 3 components: ,
- a hopf link: ,
- a Whitehead link: ,
- Borromean link: .

**Definition 1.5**



A link diagram  $D_\pi$  is a picture over projection  $\pi$  of a link  $L$  in  $\mathbb{R}^3(S^3)$  to  $\mathbb{R}^2(S^2)$  such that:

- (1)  $D_\pi|_L$  is non degenerate: ,
- (2) the double points are not degenerate: ,
- (3) there are no triple point: .

There are under- and overcrossings (tunnels and bridges) on a link diagrams with an obvious meaning.


Every link admits a link diagram.

Let  $D$  be a diagram of an oriented link (to each component of a link we add an arrow in the diagram).

We can distinguish two types of crossings: right-handed () , called a positive crossing, and left-handed () , called a negative crossing.

**1.1 Reidemeister moves**

A Reidemeister move is one of the three types of operation on a link diagram as shown below:



**Theorem 1.2** (Reidemeister, 1927 )

*Two diagrams of the same link can be deformed into each other by a finite sequence of Reidemeister moves (and isotopy of the plane).*

## 1.2 Seifert surface

Let  $D$  be an oriented diagram of a link  $L$ . We change the diagram by smoothing each crossing:

$$\begin{aligned} \nearrow \searrow &\mapsto \searrow \nearrow \\ \searrow \nearrow &\mapsto \nearrow \searrow \end{aligned}$$

We smooth all the crossings, so we get a disjoint union of circles on the plane. Each circle bounds a disk in  $\mathbb{R}^3$  (we choose disks that don't intersect). For each smoothed crossing we add a twisted band: right-handed for a positive and left-handed for a negative one. We get an orientable surface  $\Sigma$  such that  $\partial\Sigma = L$ .

Note: the obtained surface isn't unique and in general doesn't need to be connected, but by taking connected sum of all components we can easily get a connected surface (i.e. we take two disconnected components and cut a




Figure 1: Constructing a Seifert surface.

disk in each of them:  $D_1$  and  $D_2$ ; now we glue both components on the boundaries:  $\partial D_1$  and  $\partial D_2$ .

**Theorem 1.3** (Seifert)

*Every link in  $S^3$  bounds a surface  $\Sigma$  that is compact, connected and orientable. Such a surface is called a Seifert surface.*

**Definition 1.6**

*The three genus  $g_3(K)$  ( $g(K)$ ) of a knot  $K$  is the minimal genus of a Seifert surface  $\Sigma$  for  $K$ .*

**Corollary 1.1**

*A knot  $K$  is trivial if and only  $g_3(K) = 0$ .*

Remark: there are knots that admit non isotopic Seifert surfaces of minimal genus (András Juhász, 2008).




Figure 2: Connecting two surfaces.




Figure 3: Genus of an orientable surface.

**Definition 1.7**


Suppose  $\alpha$  and  $\beta$  are two simple closed curves in  $\mathbb{R}^3$ . On a diagram  $L$  consider all crossings between  $\alpha$  and  $\beta$ . Let  $N_+$  be the number of positive crossings,  $N_-$  - negative. Then the linking number:  $\text{lk}(\alpha, \beta) = \frac{1}{2}(N_+ - N_-)$ .

Let  $\alpha$  and  $\beta$  be two disjoint simple cross curves in  $S^3$ . Let  $\nu(\beta)$  be a tubular neighbourhood of  $\beta$ . The linking number can be interpreted via first homology group, where  $\text{lk}(\alpha, \beta)$  is equal to evaluation of  $\alpha$  as element of first homology group of the complement of  $\beta$ :


$$\alpha \in H_1(S^3 \setminus \nu(\beta), \mathbb{Z}) \cong \mathbb{Z}.$$

### Example 1.3

- *Hopf link:*



- *$T(6, 2)$  link:*



### Fact 1.1

$$g_3(\Sigma) = \frac{1}{2}b_1(\Sigma) = \frac{1}{2} \dim_{\mathbb{R}} H_1(\Sigma, \mathbb{R}),$$


where  $b_1$  is first Betti number of  $\Sigma$ .

### 1.3 Seifert matrix

Let  $L$  be a link and  $\Sigma$  be an oriented Seifert surface for  $L$ . Choose a basis for  $H_1(\Sigma, \mathbb{Z})$  consisting of simple closed  $\alpha_1, \dots, \alpha_n$ . Let  $\alpha_1^+, \dots, \alpha_n^+$  be copies of  $\alpha_i$  lifted up off the surface (push up along a vector field normal to  $\Sigma$ ). Note that elements  $\alpha_i$  are contained in the Seifert surface while all  $\alpha_i^+$  don't intersect the surface. Let  $\text{lk}(\alpha_i, \alpha_j^+) = \{a_{ij}\}$ . Then the matrix  $S = \{a_{ij}\}_{i,j=1}^n$  is called a Seifert matrix for  $L$ . Note that by choosing a different basis we get a different matrix.

### Theorem 1.4

The Seifert matrices  $S_1$  and  $S_2$  for the same link  $L$  are  $S$ -equivalent, that is,  $S_2$  can be obtained from  $S_1$  by a sequence of following moves:



(1)  $V \rightarrow AVA^T$ , where  $A$  is a matrix with integer coefficients,

$$(2) V \rightarrow \left( \begin{array}{ccc|cc} & & & * & 0 \\ & V & & \vdots & \vdots \\ & & & * & 0 \\ \hline * & \dots & * & 0 & 0 \\ 0 & \dots & 0 & 1 & 0 \end{array} \right) \quad \text{or} \quad V \rightarrow \left( \begin{array}{ccc|cc} & & & * & 0 \\ & V & & \vdots & \vdots \\ & & & * & 0 \\ \hline * & \dots & * & 0 & 1 \\ 0 & \dots & 0 & 0 & 0 \end{array} \right)$$

(3) inverse of (2)

## Lecture 2

March 4, 2019

### Theorem 2.1

For any knot  $K \subset S^3$  there exists a connected, compact and orientable surface  $\Sigma(K)$  such that  $\partial\Sigma(K) = K$

*Proof.* ("joke")

Let  $K \in S^3$  be a knot and  $N = \nu(K)$  be its tubular neighbourhood. Because  $K$  and  $N$  are homotopy equivalent, we get:

$$H^1(S^3 \setminus N) \cong H^1(S^3 \setminus K).$$

Let us consider a long exact sequence of cohomology of a pair  $(S^3, S^3 \setminus N)$  with integer coefficients:



$$\begin{array}{ccccccc}
& & & & \mathbb{Z} & & \\
& & & & \cong & & \\
& & & & H^0(S^3) \rightarrow H^0(S^3 \setminus N) \rightarrow & & \\
\rightarrow & H^1(S^3, S^3 \setminus N) \rightarrow & H^1(S^3) \rightarrow & H^1(S^3 \setminus N) \rightarrow & & & \\
& & & & \cong & & \\
& & & & 0 & & \\
& & & & \cong & & \\
\rightarrow & H^2(S^3, S^3 \setminus N) \rightarrow & H^2(S^3) \rightarrow & H^2(S^3 \setminus N) \rightarrow & & & \\
\rightarrow & H^3(S^3, S^3 \setminus N) \rightarrow & H^3(S) \rightarrow & & 0 & & \\
& & & & \cong & & \\
& & & & \mathbb{Z} & & 
\end{array}$$

$$H^*(S^3, S^3 \setminus N) \cong H^*(N, \partial N)$$

????????????????

□

**Definition 2.1**

Let  $S$  be a Seifert matrix for a knot  $K$ . The Alexander polynomial  $\Delta_K(t)$  is a Laurent polynomial:

$$\Delta_K(t) := \det(tS - S^T) \in \mathbb{Z}[t, t^{-1}] \cong \mathbb{Z}[\mathbb{Z}]$$

**Theorem 2.2**

$\Delta_K(t)$  is well defined up to multiplication by  $\pm t^k$ , for  $k \in \mathbb{Z}$ .

*Proof.* We need to show that  $\Delta_K(t)$  doesn't depend on  $S$ -equivalence relation.

- (1) Suppose  $S' = CSC^T$ ,  $C \in \text{GL}(n, \mathbb{Z})$  (matrices invertible over  $\mathbb{Z}$ ). Then  $\det C = 1$  and:

$$\begin{aligned}
\det(tS' - S'^T) &= \det(tCSC^T - (CSC^T)^T) = \\
&= \det(tCSC^T - CS^T C^T) = \det C \det(tS - S^T) \det C^T = \det(tS - S^T)
\end{aligned}$$

(2) Let

$$A := t \left( \begin{array}{ccc|cc} & & & * & 0 \\ & & & \vdots & \vdots \\ & S & & * & 0 \\ \hline * & \dots & * & 0 & 0 \\ 0 & \dots & 0 & 1 & 0 \end{array} \right) - \left( \begin{array}{ccc|cc} & & & * & 0 \\ & & & \vdots & \vdots \\ & S^T & & * & 0 \\ \hline * & \dots & * & 0 & 1 \\ 0 & \dots & 0 & 0 & 0 \end{array} \right) = \left( \begin{array}{ccc|cc} & & & * & 0 \\ & & & \vdots & \vdots \\ & tS - S^T & & * & 0 \\ \hline * & \dots & * & 0 & -1 \\ 0 & \dots & 0 & t & 0 \end{array} \right)$$

Using the Laplace expansion we get  $\det A = \pm t \det(tS - S^T)$ .

□

**Example 2.1**

If  $K$  is a trefoil then we can take  $S = \begin{pmatrix} -1 & -1 \\ 0 & -1 \end{pmatrix}$ . Then

$$\Delta_K(t) = \det \begin{pmatrix} -t+1 & -t \\ 1 & -t+1 \end{pmatrix} = (t-1)^2 + t = t^2 - t + 1 \neq 1 \Rightarrow \text{trefoil is not trivial.}$$

**Fact 2.1**

$\Delta_K(t)$  is symmetric.

*Proof.* Let  $S$  be an  $n \times n$  matrix.

$$\begin{aligned} \Delta_K(t^{-1}) &= \det(t^{-1}S - S^T) = (-t)^{-n} \det(tS^T - S) = \\ &= (-t)^{-n} \det(tS - S^T) = (-t)^{-n} \Delta_K(t) \end{aligned}$$

If  $K$  is a knot, then  $n$  is necessarily even, and so  $\Delta_K(t^{-1}) = t^{-n} \Delta_K(t)$ . □


**Lemma 2.1**

$$\frac{1}{2} \deg \Delta_K(t) \leq g_3(K), \text{ where } \deg(a_n t^n + \dots + a_1 t^l) = k - l.$$

*Proof.* If  $\Sigma$  is a genus  $g$  - Seifert surface for  $K$  then  $H_1(\Sigma) = \mathbb{Z}^{2g}$ , so  $S$  is an  $2g \times 2g$  matrix. Therefore  $\det(tS - S^T)$  is a polynomial of degree at most  $2g$ . □

**Example 2.2**

There are not trivial knots with Alexander polynomial equal 1, for example:



$$\Delta_{11n34} \equiv 1.$$

**Lemma 2.2** (Dehn)

Let  $M$  be a 3-manifold and  $D^2 \xrightarrow{f} M^3$  be a map of a disk such that  $f|_{\partial D^2}$  is an embedding. Then there exists an embedding  $D^2 \xrightarrow{g} M$  such that:

$$g|_{\partial D^2} = f|_{\partial D^2}.$$

**Lecture 3**

**Example 3.1**

$F : \mathbb{C}^2 \rightarrow \mathbb{C}$  a polynomial

$$F(0) = 0$$

??????????????

as a corollary we see that  $K_T^n$ , ????

is not slice unless  $m = 0$ .

**Theorem 3.1**

The map  $j : \mathcal{C} \rightarrow \mathbb{Z}^\infty$  is a surjection that maps  $K_n$  to a linear independent set. Moreover  $\mathcal{C} \cong \mathbb{Z}$

**Fact 3.1** (Milnor Singular Points of Complex Hypersurfaces)

An oriented knot is called negative amphichiral if the mirror image  $m(K)$  of  $K$  is equivalent to the reverse knot of  $K$ :  $K^r$ .

**Problem 3.1**

*Prove that if  $K$  is negative amphichiral, then  $K\#K = 0$  in  $\mathcal{C}$ .*

**Example 3.2**

*Figure 8 knot is negative amphichiral.*

## Lecture 4 Concordance group


March 18, 2019

**Definition 4.1**

*A knot  $K$  is called (smoothly) slice if  $K$  is smoothly concordant to an unknot. A knot  $K$  is smoothly slice if and only if  $K$  bounds a smoothly embedded disk in  $B^4$ .*

**Definition 4.2**

*Two knots  $K$  and  $K'$  are called (smoothly) concordant if there exists an annulus  $A$  that is smoothly embedded in  $S^3 \times [0, 1]$  such that  $\partial A = K' \times \{1\} \sqcup K \times \{0\}$ .*



Let  $m(K)$  denote a mirror image of a knot  $K$ .

**Fact 4.1**

For any  $K$ ,  $K \# m(K)$  is slice.

**Fact 4.2**

Concordance is an equivalence relation.

**Fact 4.3**

If  $K_1 \sim K_1'$  and  $K_2 \sim K_2'$ , then  $K_1 \# K_2 \sim K_1' \# K_2'$ .




Figure 4: Sketch for Fakt 4.3.

**Fact 4.4**

$K \# m(K) \sim$  the unknot.

**Theorem 4.1**

Let  $\mathcal{C}$  denote a set of all equivalent classes for knots and  $\{0\}$  denote class of all knots concordant to a trivial knot.  $\mathcal{C}$  is a group under taking connected sums. The neutral element in the group is  $\{0\}$  and the inverse element of an element  $\{K\} \in \mathcal{C}$  is  $-\{K\} = \{mK\}$ .

**Fact 4.5**

The figure eight knot is a torsion element in  $\mathcal{C}$  ( $2K \sim$  the unknot).

**Problem 4.1** (open)

Are there in concordance group torsion elements that are not 2 torsion elements?

Remark:  $K \sim K' \Leftrightarrow K \# -K'$  is slice.

Let  $\Omega$  be an oriented

???????

Suppose  $\Sigma$  is a Seifert matrix with an intersection form  $(\alpha, \beta) \mapsto \text{lk}(\alpha, \beta^+)$ .

Suppose  $\alpha, \beta \in H_1(\Sigma, \mathbb{Z})$  (i.e. there are cycles).

????????????????

$\alpha, \beta \in \ker(H_1(\Sigma, \mathbb{Z}) \rightarrow H_1(\Omega, \mathbb{Z}))$ . Then there are two cycles  $A, B \in \Omega$  such that  $\partial A = \alpha$  and  $\partial B = \beta$ . Let  $B^+$  be a push off of  $B$  in the positive normal direction such that  $\partial B^+ = \beta^+$ . Then  $\text{lk}(\alpha, \beta^+) = A \cdot B^+$

## Lecture 5

April 8, 2019

$X$  is a closed orientable four-manifold. Assume  $\pi_1(X) = 0$  (it is not needed to define the intersection form). In particular  $H_1(X) = 0$ .  $H_2$  is free (exercise).

$$H_2(X, \mathbb{Z}) \xrightarrow{\text{Poincaré duality}} H^2(X, \mathbb{Z}) \xrightarrow{\text{evaluation}} \text{Hom}(H_2(X, \mathbb{Z}), \mathbb{Z})$$

Intersection form:  $H_2(X, \mathbb{Z}) \times H_2(X, \mathbb{Z}) \rightarrow \mathbb{Z}$  - symmetric, non singular.  
Let  $A$  and  $B$  be closed, oriented surfaces in  $X$ .

### Proposition 5.1

$A \cdot B$  doesn't depend of choice of  $A$  and  $B$  in their homology classes.

Lecture 6

March 11, 2019

Definition 6.1

A link  $L$  is fibered if there exists a map  $\phi : S^3 \setminus L \leftarrow S^1$  which is locally trivial fibration.

Lecture 7

April 15, 2019

In other words:

Choose a basis  $(b_1, \dots, b_i)$

???

of  $H_2(Y, \mathbb{Z})$ , then  $A = (b_i, b_y)$

??

is a matrix of intersection form:

$$\mathbb{Z}^n / A\mathbb{Z}^n \cong H_1(Y, \mathbb{Z}).$$

In particular  $|\det A| = \#H_1(Y, \mathbb{Z})$ .

That means - what is happening on boundary is a measure of degeneracy.

$$\begin{array}{ccc}
H_1(Y, \mathbb{Z}) & \times & H_1(Y, \mathbb{Z}) & \longrightarrow & \mathbb{Q} / \mathbb{Z} & \text{- a linking form} \\
\cong & & \cong & & & \\
\mathbb{Z}^n / A\mathbb{Z} & & \mathbb{Z}^n / A\mathbb{Z} & & & 
\end{array}$$

$$(a, b) \mapsto aA^{-1}b^T$$

The intersection form on a four-manifold determines the linking on the boundary.

Let  $K \in S^1$  be a knot,  $\Sigma(K)$  its double branched cover. If  $V$  is a Seifert matrix for  $K$ , then  $H_1(\Sigma(K), \mathbb{Z}) \cong \mathbb{Z}^n / AV$  where  $A = V \times V^T$ ,  $n = \text{rank } V$ .




Figure 5: Pushing the Seifert surface in 4-ball.

Let  $X$  be the four-manifold obtained via the double branched cover of  $B^4$  branched along  $\tilde{\Sigma}$ .

**Fact 7.1**

- $X$  is a smooth four-manifold,
- $H_1(X, \mathbb{Z}) = 0$ ,
- $H_2(X, \mathbb{Z}) \cong \mathbb{Z}^n$
- The intersection form on  $X$  is  $V + V^T$ .

Let  $Y = \Sigma(K)$ . Then:

$$\begin{aligned}
 H_1(Y, \mathbb{Z}) \times H_1(Y, \mathbb{Z}) &\longrightarrow \mathbb{Q} / \mathbb{Z} \\
 (a, b) &\mapsto aA^{-1}b^T, \quad A = V + V^T \\
 H_1(Y, \mathbb{Z}) &\cong \mathbb{Z}^n / AZ \\
 A &\longrightarrow BAC^T \quad \text{Smith normal form}
 \end{aligned}$$

????????????????????????????????

In general

**Lecture 8**

May 20, 2019

Let  $M$  be compact, oriented, connected four-dimensional manifold. If  $H_1(M, \mathbb{Z}) = 0$  then there exists a bilinear form - the intersection form on  $M$ :



$$\begin{array}{c}
H_2(M, \mathbb{Z}) \quad \times \quad H_2(M, \mathbb{Z}) \longrightarrow \mathbb{Z} \\
\Downarrow \\
\mathbb{Z}^n
\end{array}$$

Let us consider a specific case:  $M$  has a boundary  $Y = \partial M$ . Betti number  $b_1(Y) = 0$ ,  $H_1(Y, \mathbb{Z})$  is finite. Then the intersection form can be degenerated in the sense that:

$$\begin{array}{ccc}
H_2(M, \mathbb{Z}) \times H_2(M, \mathbb{Z}) \longrightarrow \mathbb{Z} & & H_2(M, \mathbb{Z}) \longrightarrow \text{Hom}(H_2(M, \mathbb{Z}), \mathbb{Z}) \\
(a, b) \mapsto \mathbb{Z} & & a \mapsto (a, \_ )H_2(M, \mathbb{Z})
\end{array}$$

has coker precisely  $H_1(Y, \mathbb{Z})$ .

????????????????

Let  $K \subset S^3$  be a knot,

$X = S^3 \setminus K$  - a knot complement,

$\tilde{X} \xrightarrow{\rho} X$  - an infinite cyclic cover (universal abelian cover).

$$\pi_1(X) \longrightarrow \pi_1(X) / [\pi_1(X), \pi_1(X)] = H_1(X, \mathbb{Z}) \cong \mathbb{Z}$$

$C_*(\tilde{X})$  has a structure of a  $\mathbb{Z}[t, t^{-1}] \cong \mathbb{Z}[\mathbb{Z}]$  module.

$H_1(\tilde{X}, \mathbb{Z}[t, t^{-1}])$  - Alexander module,

$$H_1(\tilde{X}, \mathbb{Z}[t, t^{-1}]) \times H_1(\tilde{X}, \mathbb{Z}[t, t^{-1}]) \longrightarrow \mathbb{Q} / \mathbb{Z}[t, t^{-1}]$$

**Fact 8.1**

$$H_1(\tilde{X}, \mathbb{Z}[t, t^{-1}]) \cong \mathbb{Z}[t, t^{-1}]^n / (tV - V^T)\mathbb{Z}[t, t^{-1}]^n,$$

where  $V$  is a Seifert matrix.

**Fact 8.2**

$$\begin{array}{c}
H_1(\tilde{X}, \mathbb{Z}[t, t^{-1}]) \times H_1(\tilde{X}, \mathbb{Z}[t, t^{-1}]) \longrightarrow \mathbb{Q} / \mathbb{Z}[t, t^{-1}] \\
(\alpha, \beta) \mapsto \alpha^{-1}(t-1)(tV - V^T)^{-1}\beta
\end{array}$$

Note that  $\mathbb{Z}$  is not PID. Therefore we don't have primer decomposition of this moduli. We can simplify this problem by replacing  $\mathbb{Z}$  by  $\mathbb{R}$ . We lose some date by doing this transition.

$$\begin{aligned} \xi \in S^1 \setminus \{\pm 1\} \quad p_\xi &= (t - \xi)(t - \xi^{-1})t^{-1} \\ \xi \in \mathbb{R} \setminus \{\pm 1\} \quad q_\xi &= (t - \xi)(t - \xi^{-1})t^{-1} \\ \xi \notin \mathbb{R} \cup S^1 \quad q_\xi &= (t - \xi)(t - \bar{\xi})(t - \xi^{-1})(t - \bar{\xi}^{-1})t^{-2} \\ \Lambda &= \mathbb{R}[t, t^{-1}] \end{aligned}$$

$$\text{Then: } H_1(\tilde{X}, \Lambda) \cong \bigoplus_{\substack{\xi \in S^1 \setminus \{\pm 1\} \\ k \geq 0}} (\Lambda / p_\xi^k)^{n_{k, \xi}} \oplus \bigoplus_{\substack{\xi \notin S^1 \\ l \geq 0}} (\Lambda / q_\xi^l)^{n_{l, \xi}}$$

We can make this composition orthogonal with respect to the Blanchfield paring.

Historical remark:

- John Milnor, *On isometries of inner product spaces*, 1969,
- Walter Neumann, *Invariants of plane curve singularities* , 1983,
- András Némethi, *The real Seifert form and the spectral pairs of isolated hypersurfaceenumerate singularities*, 1995,
- Maciej Borodzik, Stefan Friedl *The unknotting number and classical invariants II*, 2014.

Let  $p = p_\xi$ ,  $k \geq 0$ .

$$\Lambda / p^k \Lambda \times \Lambda / p^k \Lambda \longrightarrow \mathbb{Q}(t) / \Lambda$$

$$(1, 1) \mapsto \kappa$$

$$\text{Now: } (p^k \cdot 1, 1) \mapsto 0$$

$$p^k \kappa = 0 \in \mathbb{Q}(t) / \Lambda$$

$$\text{therefore } p^k \kappa \in \Lambda$$

$$\text{we have } (1, 1) \mapsto \frac{h}{p^k}$$

$h$  is not uniquely defined:  $h \rightarrow h + gp^k$  doesn't affect paring.  
 Let  $h = p^k \kappa$ .

**Example 8.1**

$$\begin{aligned}\phi_0((1, 1)) &= \frac{+1}{p} \\ \phi_1((1, 1)) &= \frac{-1}{p}\end{aligned}$$

$\phi_0$  and  $\phi_1$  are not isomorphic.

*Proof.* Let  $\Phi : \Lambda/p^k\Lambda \rightarrow \Lambda/p^k\Lambda$  be an isomorphism.  
 Let:  $\Phi(1) = g \in \lambda$

$$\begin{aligned}\Lambda/p^k\Lambda &\xrightarrow{\Phi} \Lambda/p^k\Lambda \\ \phi_0((1, 1)) = \frac{1}{p^k} &\quad \phi_1((g, g)) = \frac{1}{p^k} \quad (\Phi \text{ is an isometry}).\end{aligned}$$

Suppose for the paring  $\phi_1((g, g)) = \frac{1}{p^k}$  we have  $\phi_1((1, 1)) = \frac{-1}{p^k}$ . Then:

$$\begin{aligned}\frac{-g\bar{g}}{p^k} &= \frac{1}{p^k} \in \mathbb{Q}(t)/\Lambda \\ \frac{-g\bar{g}}{p^k} - \frac{1}{p^k} &\in \Lambda \\ -g\bar{g} &\equiv 1 \pmod{p} \text{ in } \Lambda \\ -g\bar{g} - 1 &= p^k\omega \text{ for some } \omega \in \Lambda\end{aligned}$$

evaluating at  $\xi$ :

$$\overbrace{-g(\xi)g(\xi^{-1})}^{>0} - 1 = 0 \quad \Rightarrow \Leftarrow$$

□

????????????????????

$$\begin{aligned}
g &= \sum g_i t^i \\
\bar{g} &= \sum g_i t^{-i} \\
\bar{g}(\xi) &= \sum g_i \xi^i \quad \xi \in S^1 \\
\bar{g}(\xi) &= g(\bar{\xi})
\end{aligned}$$

Suppose  $g = (t - \xi)^\alpha g'$ . Then  $(t - \xi)^{k-\alpha}$  goes to 0 in  $\Lambda/p^k\Lambda$ .

**Theorem 8.1**

*Every sesquilinear non-degenerate pairing*

$$\Lambda/p^k \times \Lambda/p \leftrightarrow \frac{h}{p^k}$$

*is isomorphic either to the pairing with  $h = 1$  or to the pairing with  $h = -1$  depending on sign of  $h(\xi)$  (which is a real number).*

*Proof.* There are two steps of the proof:

1. Reduce to the case when  $h$  has a constant sign on  $S^1$ .
2. Prove in the case, when  $h$  has a constant sign on  $S^1$ .

**Lemma 8.1**

*If  $P$  is a symmetric polynomial such that  $P(\eta) \geq 0$  for all  $\eta \in S^1$ , then  $P$  can be written as a product  $P = g\bar{g}$  for some polynomial  $g$ .*

*Sketch of proof.* Induction over  $\deg P$ .

Let  $\zeta \notin S^1$  be a root of  $P$ ,  $P \in \mathbb{R}[t, t^{-1}]$ . Assume  $\zeta \notin \mathbb{R}$ . We know that polynomial  $P$  is divisible by  $(t - \zeta)$ ,  $(t - \bar{\zeta})$ ,  $(t^{-1} - \zeta)$  and  $(t^{-1} - \bar{\zeta})$ . Therefore:

$$\begin{aligned}
P' &= \frac{P}{(t - \zeta)(t - \bar{\zeta})(t^{-1} - \zeta)(t^{-1} - \bar{\zeta})} \\
P' &= g'\bar{g}
\end{aligned}$$

We set  $g = g'(t - \zeta)(t - \bar{\zeta})$  and  $P = g\bar{g}$ . Suppose  $\zeta \in S^1$ . Then  $(t - \zeta)^2 \mid P$  (at least - otherwise it would change sign). Therefore:

$$\begin{aligned}
P' &= \frac{P}{(t - \zeta)^2(t^{-1} - \zeta)^2} \\
g &= (t - \zeta)(t^{-1} - \zeta)g' \quad \text{etc.}
\end{aligned}$$

The map  $(1, 1) \mapsto \frac{h}{p^k} = \frac{g\bar{g}h}{p^k}$  is isometric whenever  $g$  is coprime with  $P$ .  $\square$

**Lemma 8.2**

Suppose  $A$  and  $B$  are two symmetric polynomials that are coprime and that  $\forall z \in S^1$  either  $A(z) > 0$  or  $B(z) > 0$ . Then there exist symmetric polynomials  $P, Q$  such that  $P(z), Q(z) > 0$  for  $z \in S^1$  and  $PA + QB \equiv 1$ .

*Idea of proof.* For any  $z$  find an interval  $(a_z, b_z)$  such that if  $P(z) \in (a_z, b_z)$  and  $P(z)A(z) + Q(z)B(z) = 1$ , then  $Q(z) > 0$ ,  $x(z) = \frac{az+bz}{i}$  is a continuous function on  $S^1$  approximating  $z$  by a polynomial .  
 ??

$$(1, 1) \mapsto \frac{h}{p^k} \mapsto \frac{g\bar{g}h}{p^k}$$

$$g\bar{g}h + p^k\omega = 1$$

Apply Lemma 8.2 for  $A = h, B = p^{2k}$ . Then, if the assumptions are satisfied,

$$Ph + Qp^{2k} = 1$$

$$p > 0 \Rightarrow p = g\bar{g}$$

$$p = (t - \xi)(t - \bar{\xi})t^{-1}$$

$$\text{so } p \geq 0 \text{ on } S^1$$

$$p(t) = 0 \Leftrightarrow t = \xi \text{ or } t = \bar{\xi}$$

$$h(\xi) > 0$$

$$h(\bar{\xi}) > 0$$

$$g\bar{g}h + Qp^{2k} = 1$$

$$g\bar{g}h \equiv 1 \pmod{p^{2k}}$$

$$g\bar{g} \equiv 1 \pmod{p^k}$$

??  
 If  $P$  has no roots on  $S^1$  then  $B(z) > 0$  for all  $z$ , so the assumptions of Lemma 8.2 are satisfied no matter what  $A$  is. □

????????????????????????????

$$\begin{aligned}
(\Lambda/p_\xi^k \times \Lambda/p_\xi^k) &\longrightarrow \frac{\epsilon}{p_\xi^k}, & \xi \in S^1 \setminus \{\pm 1\} \\
(\Lambda/q_\xi^k \times \Lambda/q_\xi^k) &\longrightarrow \frac{1}{q_\xi^k}, & \xi \notin S^1
\end{aligned}$$

???????????????????? 1 ?? epsilon?

**Theorem 8.2**

(Matumoto, Conway-Borodzik-Politarczyk) Let  $K$  be a knot,

$$H_1(\tilde{X}, \Lambda) \times H_1(\tilde{X}, \Lambda) = \bigoplus_{\substack{k, \xi, \epsilon \\ \xi \in S^1}} (\Lambda/p_\xi^k, \epsilon)^{n_{k, \xi, \epsilon}} \oplus \bigoplus_{k, \eta} (\Lambda/p_\xi^k)^{m_k}$$

$$\text{Let } \delta_\sigma(\xi) = \lim_{\epsilon \rightarrow 0^+} \sigma(e^{2\pi i \epsilon \xi}) - \sigma(e^{-2\pi i \epsilon \xi}),$$

$$\text{then } \sigma_j(\xi) = \sigma(\xi) - \frac{1}{2} \lim_{\epsilon \rightarrow 0} \sigma(e^{2\pi i \epsilon \xi}) + \sigma(e^{-2\pi i \epsilon \xi})$$

The jump at  $\xi$  is equal to  $2 \sum_{k_i \text{ odd}} \epsilon_i$ . The peak of the signature function is equal to  $\sum_{k_i \text{ even}} \epsilon_i$ .

□

**Lecture 9**

May 27, 2019

....

**Definition 9.1**

A square hermitian matrix  $A$  of size  $n$ .

field of fractions

**Theorem 10.1**

Let  $K$  be a knot and  $u(K)$  its unknotting number. Let  $g_4(K)$  be a minimal four genus of a smooth surface  $S$  in  $B^4$  such that  $\partial S = K$ . Then:

$$u(K) \geq g_4(K)$$

*Proof.* Recall that if  $u(K) = u$  then  $K$  bounds a disk  $\Delta$  with  $u$  ordinary double points.

Remove from  $\Delta$  the two self intersecting and glue the Seifert surface for the Hopf link. The reality surface  $S$  has Euler characteristic  $\chi(S) = 1 - 2u$ . Therefore  $g_4(S) = u$ . □

????????????????????

**Example 10.1**

The knot  $8_{20}$  is slice:  $\sigma \equiv 0$  almost everywhere but  $\sigma(e^{\frac{2\pi i}{6}}) = +1$ .

**Surgery**

Recall that  $H_1(S^1 \times S^1, \mathbb{Z}) = \mathbb{Z}^2$ . As generators for  $H_1$  we can set  $\alpha = [S^1 \times \{pt\}]$  and  $\beta = [\{pt\} \times S^1]$ . Suppose  $\phi : S^1 \times S^1 \rightarrow S^1 \times S^1$  is a diffeomorphism.

Consider an induced map on homology group:

$$\begin{aligned} H_1(S^1 \times S^1, \mathbb{Z}) \ni \phi_*(\alpha) &= p\alpha + q\beta, & p, q \in \mathbb{Z}, \\ \phi_*(\beta) &= r\alpha + s\beta, & r, s \in \mathbb{Z}, \\ \phi_* &= \begin{pmatrix} p & q \\ r & s \end{pmatrix} \end{aligned}$$

As  $\phi_*$  is diffeomorphis, it must be invertible over  $\mathbb{Z}$ . Then for a direction preserving diffeomorphism we have  $\det \phi_* = 1$ . Therefore  $\phi_* \in \text{SL}(2, \mathbb{Z})$ .

## Lecture 11 balagan

*Proof.* By Poincaré duality we know that:

$$\begin{aligned} H_3(\Omega, Y) &\cong H^0(\Omega), \\ H_2(Y) &\cong H^0(Y), \\ H_2(\Omega) &\cong H^1(\Omega, Y), \\ H_2(\Omega, Y) &\cong H^1(\Omega). \end{aligned}$$

Therefore  $\dim_{\mathbb{Q}} H_1(Y)/_V = \dim_{\mathbb{Q}} V$ . □

Suppose  $g(K) = 0$  ( $K$  is slice). Then  $H_1(\Sigma, \mathbb{Z}) \cong H_1(Y, \mathbb{Z})$ . Let  $g_{\Sigma}$  be the genus of  $\Sigma$ ,  $\dim H_1(Y, \mathbb{Z}) = 2g_{\Sigma}$ . Then the Seifert form  $V$  on a 4 - manifolds???

?????

has a subspace of dimension  $g_{\Sigma}$  on which it is zero:

$$V = \left\{ \begin{array}{c} \overbrace{\left( \begin{array}{cccccc} 0 & \dots & 0 & * & \dots & * \\ \vdots & & \vdots & \vdots & & \vdots \\ 0 & \dots & 0 & * & \dots & * \\ * & \dots & * & * & \dots & * \\ \vdots & & \vdots & \vdots & & \vdots \\ * & \dots & * & * & \dots & * \end{array} \right)}^{g_{\Sigma}} \\ \end{array} \right\}_{2g_{\Sigma} \times 2g_{\Sigma}}$$